

ASSESSMENT OF ICHTHYOFAUNAL DIVERSITY AND PHYSICOCHEMICAL PARAMETERS IN BHADAR-II RESERVOIR, GUJARAT

Abstract

The study was conducted to assess the fish diversity and water quality in the Bhadar-II reservoir, located in Rajkot district of Gujarat. Monthly sampling was carried out from September 2023 to April 2024. A total of 19 fish species were recorded during the study period. Cypriniformes was the dominant order in terms of species diversity (9 species), followed by Siluriformes (5 species), Anabantiformes (3 species), Osteoglossiformes (1 species) and Cichliformes (1 species). The dominant families were Cyprinidae (9 species), followed by Channidae (3 species), Siluridae (2 species), and 1 species each from Claridae, Heteropneustidae, Cichlidae, Pangasiidae, and Notopteridae. The catfish population was the most prevalent, accounting for 26% of the total fish population, followed by minor carp at 21%. Major carp, exotic fish, and murrels each contributed 16%, while featherback fish had the lowest at 5%. The fish data was subjected to species diversity analysis. The Shannon-Wiener Index values ranged from 1.776 to 2.546, indicating monthly fluctuations in diversity levels. This range suggests that the habitat structure was moderately stable. Pielou's Evenness Index values ranged from 0.8080 to 0.8880, showing a gradual increase in evenness over time. This increase can be attributed to a decline in the population of the dominant species. Species richness, as measured by the Margalef Richness Index, ranged from 1.278 to 2.477, with the highest species richness observed in November during the winter season. Monthly observations of physicochemical parameters showed the range of water temperature from 23.4°C to 30.1°C, dissolved oxygen levels from 6.7 to 8.9 mg/l, pH from 7.6 to 8.3, alkalinity between 139.4 to 156.2 mg/l, and total hardness from 94.2 to 133.1 mg/l.

Keywords: Ichthyofaunal, Diversity Indices, Water Quality, Reservoir.

1. Introduction

Biodiversity, encompassing species, ecosystem, and genetic diversity, reflects the vast variety of life forms on earth, with uneven distribution favouring tropical regions (Walker, 1992; Jana et al., 2015). Ichthyofaunal diversity, referring to the variety of fish in aquatic ecosystems, is crucial for ecological balance and food security. Globally, fish comprise over 50% of vertebrate species, with India contributing 7.7% of this diversity, including 1,673

marine and 994 freshwater species (Froese & Pauly, 2020). Freshwater ecosystems, covering 0.8% of India's land, support 9.7% of its fauna and provide key ecological services like flood control and water purification (Pathak & Lavudya, 2021).

Reservoirs, a vital component of India's inland fisheries, span 3.15 million hectares across 19,370 water bodies (Desai, 2006). These reservoirs are classified into small, medium, and large categories, with small reservoirs playing a critical role in fisheries, offering higher yields. Despite their potential, reservoir fisheries face challenges such as poor management, overexploitation, and habitat disruption. Gujarat, home to 711 reservoirs, has significant aquaculture potential, although its fisheries are underdeveloped (Sugunan, 1995). The diversity of ichthyofauna within these ecosystems is vital for sustenance, economic benefits, and biodiversity. However, threats like pollution, overfishing, and river modifications jeopardize this diversity, with 20% of global freshwater fish species at risk (Postle, 2002). Advancing sustainable management practices is imperative to harness the full potential of India's reservoirs and preserve ichthyofaunal diversity.

2. Materials and methods

2.1 Sampling site

The research was carried out in the Saurashtra region of Gujarat, India, specifically at the Bhadar-II reservoir (21°45'40"N, 70°25'22"E), located in Bhukhi village, Rajkot district, approximately 120 km from the College of Fisheries Science, Veraval. The Bhadar River, located in the Kathiawar (Saurashtra) peninsula, originates near Vaddi (Aniali Village), Rajkot district, Gujarat. The Bhadar-II reservoir, a medium reservoir with a capacity of 49 million cubic meters, plays a significant role in the region. Constructed in 2000, it covers an area of 850 hectares with an effective storage capacity of 41.85 million cubic meters and experiences an average annual rainfall of 586 mm.

2.2 Sampling methods

Data collection was conducted at the study location where water temperature was measured using a digital thermometer, and pH levels were determined in situ using a digital pH meter. Water samples were then carefully transported to the laboratory for further analysis. Dissolved oxygen (DO) levels were collected in glass stopper bottles to prevent exposure to air. Laboratory analysis of water samples included determining parameters such as total hardness (TH), dissolved oxygen (DO), and total alkalinity (TA), following standard

laboratory protocols as outlined by the American Public Health Association (APHA, 2005). Fish specimens were randomly collected using 8-12 gill nets, each measuring 8-10 meters in length and 3-5 meters in width, with mesh sizes ranging from 6 to 9 cm. These nets were deployed for 14 to 16 hours. After collection, fish specimens were identified and photographed on-site before being transported to the College of Fisheries Science, Veraval, for further examination, measurement, and identification. Upon arrival at the lab, the specimens were preserved. Fish samples were preserved using a 10% formalin solution (40% commercial solution) in separate containers based on size. Small fish were directly submerged, while larger specimens were dissected before preservation. Species identification and confirmation were conducted using available literature (Day, 1878; Talwar & Jhingran, 1991; Jayaram, 1999). Valid species nomenclature was referenced from the Catalogue of Fishes of the California Academy of Sciences (Eschmeyer & Fricke, 2024), and fish conservation status was cross-verified using the IUCN Red List (IUCN, 2024). Fish species diversity was analyzed using various indices, including species richness (Margalef, 1958), Shannon-Wiener diversity index (1963), and species evenness index (Pielou's, 1977), analysis was performed using freely available statistical software, PAST.

3. Result and Discussion

A total of 19 fish species were documented (Plate1), categorized into 5 orders (Cypriniformes, Siluriformes, Cichliformes, Anabantiformes, and Osteoglossiformes), 8 families, and 13 genera (Table.1). The dominant order was Cypriniformes, accounting for 48% of the total species, with 9 species such as *Puntius sarana*, *Labeocatla*, *Labeorohita*, *Labeoboggut*, *Labeofimbriatus*, *Labeocalbasu*, *Ctenopharyngodonidella*, and *Cyprinus carpio* and *Cirrhinus mrigala*. Other orders observed included Siluriformes (26%), Anabantiformes (16%), and Osteoglossiformes and Cichliformes (5% each) (Figure.1). The Cyprinidae family was most prevalent, aligning with Vyas et al. (2020), who also recorded a strong presence of Cyprinidae in their study. Other families, such as Channidae (3 species), Siluridae (2 species), and Cichlidae (1 species), contributed to the ichthyofaunal diversity (Figure.2).

Comparative studies reveal variation in species composition across regions. For instance, Battul et al. (2007) documented 18 species in Maharashtra, while Raja et al. (2015) recorded 48 species in Tamil Nadu, highlighting geographical and hydrological influences. The dominance of Cypriniformes is consistent with findings by Rao & Rao (2023), who observed

a similar trend in Andhra Pradesh, whereas Niyazi et al. (2020) reported even higher dominance in Chhattisgarh. Environmental factors, including water quality and physico-chemical conditions, play a significant role in species distribution and diversity. Stressful conditions reduce diversity, often leading to dominance by a few species (Plafkin et al., 1989). This study emphasizes the unique characteristics and biodiversity of the reservoir, contributing to the understanding of regional ichthyofaunal patterns.




Figure 1.Order-wise species composition of fishdiversity of Bhadar-II reservoir

Figure 2. Family-wise species composition of fish diversity of Bhadar -II reservoir

The indices of fish species were calculated based on the abundance of species of fish diversity. The Shannon-Wiener diversity index (H') for fish in the Bhadar-II reservoir varied monthly. It peaked in December at 2.546 and dropped to 1.776 in September. Higher diversity in December was likely due to favorable environmental conditions, while lower diversity in September could reflect post-monsoon changes. These results align with Sharma et al. (2020), who reported similar seasonal trends in the Dilawara reservoir. Monthly variations in species evenness, measured by Pielou's Evenness Index (J'), ranged from 0.8080 to 0.8880. The highest evenness occurred in April, and the lowest in September. Higher evenness in April was associated with a decline in dominant species populations. The findings are consistent with Raja et al. (2015), who reported similar values from the Krishnagiri Reservoir. The Margalef Richness Index (d) for fish ranged from 1.278 in September to 2.477 in November, with November showing the highest species richness due to optimal environmental conditions. Conversely, September's lower richness might be attributed to reduced post-monsoon activities. These findings are comparable to Mishra et al. (2014), though the biodiversity was lower than typically observed in healthy ecosystems (Table 1).

Table 1. Monthly variation in species diversity indices

Index	Months							
	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.
Shannon-Wiener Diversity (H')	1.776	2.238	2.523	2.546	2.477	2.415	2.303	2.278
Pielou's Evenness (J')	0.8082	0.8480	0.8569	0.8646	0.8744	0.8711	0.8728	0.8880
Margalef Richness (d)	1.278	1.862	2.477	2.437	2.167	2.064	1.838	1.763

Water quality in India shows regional variation due to diverse geological and geographical factors. This study assessed key parameter- temperature, dissolved oxygen (DO), pH, alkalinity, and total hardness, using standard analytical methods. Water temperature, influenced by collection timing, season, and depth, directly affects aquatic ecosystems. The maximum temperature recorded was 30.1°C in April (summer), and the minimum was 23.4°C in December (winter). Similar patterns were observed by Chandravanshi et al. (2020) in the Hiran-II reservoir, Gujarat, and Lianthuamluaia et al. (2013) in the Savitri reservoir, Maharashtra. DO levels were highest in December (8.9 mg/L) and lowest in April (6.7 mg/L), reflecting seasonal variations. Winter solubility increases DO, while summer reduces it. Comparable trends were noted by Chaudhari et al. (2015) at Hatnur Reservoir, Maharashtra. The pH ranged from 7.6 in December to 8.3 in April, aligning with medium productivity (Jhingran et al., 1990). Summer increases in pH were due to higher photosynthetic activity. Similar findings were reported by Vyas (2020) in the Ozat-II reservoir. Alkalinity varied between 139.4 mg/L in December and 156.2 mg/L in April, influenced by runoff from agricultural areas. These results are consistent with Vyas (2020) and Verma et al. (2011). Total hardness ranged from 94.2 mg/L in December to 133.1 mg/L in April, categorizing the water as moderately hard (Kannan, 1991). Higher hardness during summer correlates with evaporation. This aligns with Vyas et al. (2019) (Table 2).

4. Conclusion

This study investigated the monthly variations in fish diversity and water quality parameters (dissolved oxygen, temperature, alkalinity, pH, and total hardness) in the Bhadar-II reservoir, Gujarat. The recorded water quality parameters were found within the permissible limits of

BIS standards. A total of 19 fish species were documented, classified into five orders, eight families, and 13 genera. The catfish population was the most prevalent, accounting for 26% of the total fish population, followed by minor carp at 21%. Major carp, exotic fish, and murrels each contributed 16%, while featherback fish had the lowest at 5%. The fish community was predominantly composed of species from the Cypriniformes order, particularly the Cyprinidae family. The species diversity and richness indices indicated higher fish diversity and richness during the winter season (November, December, January, and February) followed by the summer season (March and April) and the post-monsoon season (September and October). The species evenness index revealed the most even distribution of fish species in April, due to a gradual decline of dominant species, and the least even distribution in September, due to the uneven distribution of fish species. The present study will form a baseline database for the conservation and management of fisheries resources of Bhadar-II reservoir. The findings of the present study will form a basis for future research on taxonomy, biology, and proximate composition of fishes from the Bhadar-II reservoir.

5. References

American Public Health Association. APHA. (1998). Standard methods for the examination of water and wastewater. *American Public Health Association, American Water Works Association, Water Environment Federation, Washington*, 6,1325-1368.

- Battul, P., Rao, R., Navale, K., Bagale, M., & Shah, (2007). Fish diversity from Ekruk Lake near Solapur, Maharashtra. *Journal of Aquatic Biology*. 22(2), 68-72.
- Chandravanshi, R., Mishra, V., Pargi, N., & Vaghela, D. T. (2020). Ichthyofaunal diversity of Hiran-ii reservoir, Gujarat with special reference to Physico-chemical parameters. *Journal of Pharmacognosy and Phytochemistry*, 9(3S), 72-76.
- Chaudhari, N. D., Lohar, P. S., & Patil, A. G. (2015). Correlation of Limnological Features with Ichthyofauna in Hatnur Reservoir in Jalgaon, Maharashtra, India. *SSR Institute of International Journal of Life Sciences*, 2581-8740.DOI: 10.21276/SSR-IJLS.2022.8.2.4
- Day, F. S. (1878). The fishes of India, being a Natural History of Fishes found to inhabit the Seas and Freshwater of India, Burma and Ceylon. Text and atlas in 4 parts. London, 1-778.
- Desai, V. R. (2006). Reservoir fisheries. *Handbook of fisheries and aquaculture. Directorate of Information and Publications of Agriculture, Indian Council of Agricultural Research, New Delhi*, 173-195.
- Fricke, R., Eschmeyer, W.N., & Van der Laan, R. (2024). Eschmeyer's Catalog of Fishes: Genera, Species, References, Version 05/2024, California Academy of Sciences. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>.
- Froese, R. & Pauly, D. (2020). FishBase. 2020. World Wide Web Electronic Publication. Available at: <http://www.fishbase.org> (Accessed on 8 January 2018).
- IUCN 2024. The IUCN Red List of Threatened Species. Version 2023-1. <<https://www.iucnredlist.org>> ISSN 2307-8235
- Jana, A., Sit, G., & Maiti, K. (2015). Ichthyofaunal diversity of Keleghai river at Medinipur district in West Bengal. *International Research Journal of Basic and Applied Sciences*, 1(2015), 24-26.
- Jayaram, K. C. (1999). The fresh water fishes of indian, region. *Hindustan publishing corporation (India), Dehli*, 3-55.
- Jhingran, A. G., & Sugunan, V. V. (1990). General guidelines and planning criteria for small reservoir fisheries management. In *Proceedings of the national workshop on reservoir fish*, 93(1), 1-8.

- Lianthumluaia, C. S., Deshmukhe, G., Landge, A. T., & RAMTEKE, K. K. (2013). Assessment of seasonal variations of water quality parameters of Savitri reservoir, Poladpur, Raigad district, Maharashtra. *The Bioscan*, 8(4), 1337-1342. <http://krishi.icar.gov.in/jspui/handle/123456789/53284>
- Margalef, D. R. 1958. Information theory in ecology. *General Systems*. 3, 36-71.
- Mishra, A., Chakraborty, S. K., Jaiswar, A. K., & Sharma, A. P. (2014). Comparative Biodiversity of Two Medium Reservoirs of North India. *National Academy Science Letters*, 5(37), 423-430.
- Niyazi, A., Swarnkar, S., Keer, N. R., Sahu, D., & Singh, J. (2020). Ichthyofaunal diversity of kharung reservoir in Bilaspur district, Chhattisgarh, India. *Journal of Experimental Zoology*, 23, 1317-1323.
- Pathak, S., & Lavudya, N. (2021). Diversity of freshwater fish in Narmada River, Madhya Pradesh. *Journal of Entomology and Zoology Studies*, 9(2), 704-709.
- Pielou, E. C. (1966). The measurement of diversity in different types of biological collections. *Journal of theoretical biology*, 13, 131-144.
- Plafkin, J. L., Barbour, M. T., Porter, K. D., Gross, S. K., & Hughes, R. M. (1989). Rapid bioassessment protocols for use in streams and rivers. *Benthic macroinvertebrates and fish. Office of Water Regulations and Standards, US Environmental Protection Agency, Washington, DC, EPA/444/4-89/001*.
- Postel, S. (2002). Water and sustainability: Dimensions of the global challenge, Global Water Policy Project. *World Watch Institute, Amherst, Massachusetts*, 22-75.
- Raja, M., Ramkumar, R., Kavitha, M., Balasubramanian, V., Raja, R. K., Anandhi, M., & Perumal, P. (2015). Ichthyofauna diversity in relation to water quality and habitat characteristics of Krishnagiri Reservoir, Tamil Nadu, India. *International Journal of Pure and Applied Biotechnology IJPAB*, 2, 1-15.
- Rao, R. R., & Rao, K. R. (2023). An Annotated Check List of Ichthyofaunal Diversity of Madduvalasa Reservoir at Srikakulam District, Andhra Pradesh, India. *Uttar Pradesh Journal of Zoology*, 44(22), 60-73.
- Shannon, C. E. & Weaver, W. (1963). *The Mathematical Theory of Communication*. University of Illinois Press, Urbana, Illinois, 1-117.

- Sharma, M., Wala, H., & Sharma, R. (2020). Ichthyofaunal diversity and productivity of Dilawara reservoir of Dhar district, Madhya Pradesh. *International Journal of Fisheries and Aquatic Studies*, 8(5), 375-379.
- Sugunan, V. V. (1995). Reservoir fisheries of India. FAO, Rome, Italy. *Fisheries Technical Paper*, 345.
- Talwar, P. K., & Jhingran, A. G. (1991). *Inland fishes of India and adjacent countries* (Vol. 2). CRC press.
- Verma, S. R., Chaudhari, P. R., Singh, R. K., & Wate, S. R. (2011). Studies on the ecology and trophic status of an urban lake at Nagpur city, India. *Rasayan J. Chem*, 4(3), 652-659.
- Vyas, T., Roy, S. M., & Choudhary, L. (2020). Ichthyofaunal Diversity of the Mahi Bajaj Sagar Reservoir, District Banswara, Rajasthan, India. *International Journal for Research in Applied Science & Engineering Technology*, 8(11), 371-375.
- Vyas, U. D., Desai, A. Y., & Temkar, G. S. (2019). Ichthyofaunal diversity of Bhadar-I reservoir, Gujarat. *Journal of Experimental Zoology, India*, 22(1), 55-60.
- Walker, B. H. (1992). Biological and ecological redundancy. *Conserv. Biol*, 6, 18-23.

Plate1: Fish specimens collected from the Bhadar-II reservoir



Labeocatla (WoRMS, 2023)



Labeorohita (Hamilton, 1822)



Labeocalbasu (Hamilton, 1822)



Labeoboggut (Sykes, 1839)



Cirrhinus mrigala (Hamilton, 1822)



Ctenopharyngodon idella (Valenciennes, 1844)



Cyprinus carpio (Linnaeus, 1758)



Puntius sarana
(Hamilton-Buchanan, 1822)



Labeo fimbriatus (Bloch, 1795)



Oreochromis mossambicus
(Peters, 1852)



Heteropneustes fossilis (Bloch, 1794)



Clarias batrachus (Linnaeus, 1758)



Ompok pabda (Hamilton, 1822)



Pangasius pangasius
(Hamilton, 1822)



Wallago attu (Bloch & Schneider, 1801)



Notopterus notopterus
(Pallas, 1769)



Channa punctata (Bloch, 1793)



Channa striata (Bloch, 1793)



Channa marulius (Hamilton, 1822)

UNDER PEER REVIEW