

Original Research Article

Alternative Pest Control Method for *Myzus persicae* (Sulzer) with the Application of Mineral Oils

Abstract

A component that is used to lessen the spread of Potato Virus Y in potato fields is mineral oil. Nevertheless, nothing is known regarding further impacts that oil can have on potato insect pests. We conducted a number of bioassay tests to determine the impact of mineral oil on the mortality of potato aphids, green peach aphids, and *Myzus persicae* (Sulzer) (Hemiptera: Aphididae) in order to gain a better understanding of how mineral oil impacts potato pests. After performing bioassay in laboratory conditions mortality was studied and calculated with one-way anova results. Mortality was recorded for different oil concentrations at different hours. Mean values for the average mean mortality varied from 15% to 75% depending on the mineral oil concentrations used. Aphids showed lesser movement on plants sprayed with mineral oil as compared to non-sprayed plants.

Keywords: PVY^o; mineral oil; aphids; *Myzus persicae*,

Introduction

For more than a century, mineral oil, a petroleum-based substance, has been utilized in pest control [1]. Numerous applications exist for mineral oil, most commonly as an insecticide. Although it can kill some lepidopteran larvae [2], its primary target is small, soft-bodied arthropods like aphids, mites, and scales [3,4,5]. Oil frequently results in lower mortality than synthetic pesticides, despite its potential for effectiveness [6,7]. The mode of action of mineral oil has received comparatively less study attention in comparison to other pesticide classes.

Although mineral oil sprays are frequently used on potatoes, little is known about their potential to assist manage insect pests in this crop. Potato producers will probably use less of other insecticides if oil is proven to have broad insecticidal or repellent qualities, which will save money and lessen the impact on the environment. Despite some research, the effects of mineral oil on aphids (*Hemiptera: Aphididae*) remain unclear, partly due to the apparent contradictions in some of the published findings.

The impact of mineral oil on aphids are detailed in this paper. Aphids named green peach aphid, *Myzus persicae* (Sulzer) (*Hemiptera: Aphididae*), were used in the tests. This species is among the most prevalent aphids that colonize potatoes.

Methodology

Sub culture of aphids was continuously be done on new Chinese cabbage plants. Average sized aphids were used for the experiment. The *Myzus persicae* colony was initiated from a single apterous parthenogenetic female. Aphids were reared on Chinese cabbage plants raised in a growth room maintained at $20 \pm 1^{\circ}\text{C}$ under a photoperiod of L16:D8. Young adults were used for experiments. Chinese cabbage seeds have been sown in pots and placed in the glass house. After few days, plants were ready for the bioassay experiments. Three mineral oils named servo, orchol and agrosipel with three concentration each were selected for the experiment. Three mineral oil concentrations (0.1% v/v, 0.5% v/v and 1% v/v are used for the spray. Vials were filled with agar solution and leaves were cut in a disc shape before placing

them in experimental vials. Agar solution was prepared one night before setting the experiment. Leaves were treated with different concentrations of mineral oils. Leaves were air dried before placing them in agar filled vials. After that aphid were placed on the leaves to feed and vials were covered with muslin cloth before placed in incubator in 21⁰ C temperature. Thereafter, readings were recorded till 100% mortality rate or leaves got dry completely. The experiment was repeated several times. One-way anova was performed to analyse mineral oil effects on aphid mortality. After performing anova parametric tukey test was performed using software spss. Results are reported as mean \pm standard error of the mean (SEM)

Results

1.Servo leaf treatment

Leaf discs were sprayed with servo oil concentrations and placed in incubator. Readings were recorded each hour for 5-6 days till the leaf disc got dried or aphids shown 100% mortality. Later on, normality tests were run for the values and one-way anova readings were recorded. From the table 1. it is observed that the mean and median for the treatment are almost equal and skewness for the constructs is nearer to zero, hence it confirms asymptotic normality of the data. This enables to apply parametric test on the collected sample data. Mineral oil application on leaf with different concentrations showed significant results. Mortality showing mean \pm SE as 6.19 \pm 2.63, 15.9 \pm 3.51, 30.9 \pm 4.34 and 65.7 \pm 4.34 was observed for servo (df =3, mean square =4775.126 between groups and df=24, mean square =100.857) with significance <.001. After performing post hoc test for servo, it was found out in table 3 that control, 0.1v/v, 0.5%v/v and 1% v/v are showing significant results. After leaf treatment servo showed significant mortality with each concentration. Bar graph 1 showing six days average mortality for 0.1% v/v servo 15.9%. 0.5%v/v servo showed mortality in aphids to 30.9%. Average mean mortality for 1% servo was 65.7%. it didn't harm the potato leaves in any way. After leaf treatment servo showed significant mortality with each concentration. Bar graph 1 showing six days average mortality for 0.1% v/v servo 15.9%. 0.5%v/v servo showed mortality in aphids to 30.9%. Average mean mortality for 1% servo was 65.7%. it didn't harm the potato leaves in any way.

	Mean \pm SE	Median	SD	Skewness	Kurtosis
Control		3.33		1.568	2.260
6.19 \pm 2.63		6.96			
0.1%	servo	14.2		-.001	-1.428
15.9 \pm 3.51		9.29			
0.5%	servo	33.3		-.235	-.816
30.9 \pm 4.34		11.5			
1.0%	servo	63.3		-.174	0.810
65.7 \pm 4.34		11.5			

Table 1: Normality readings for servo leaf treatment.


Percentage Mortality					
	Sum of Squares	df	Mean Square	F	Sig.
Between	14325.377	3	4775.126	47.345	<.001

Groups					
Within Groups	2420.578	24	100.857		
Total	16745.955	27			

Table 2: One-way anova result for servo concentrations

Type of treatment (I)	Type of treatment (J)	Mean difference (I-J)	significance
Control	0.1% Servo	-9.55095	<.001
	0.5% Servo	-24.76667 [*]	<.001
	1% Servo	-59.52857 [*]	<.001
0.1% Servo Mortality	Control	9.55095	<.001
	0.5% Servo	-15.21571 [*]	<.001
	1% Servo	-49.97762 [*]	<.001
0.5% Servo Mortality	Control	24.76667 [*]	<.001
	0.1% Servo	15.21571 [*]	<.001
	1% Servo	-34.76190 [*]	<.001
1% Servo Mortality	Control	59.52857 [*]	<.001
	0.1% Servo	49.97762 [*]	<.001
	0.5% Servo	34.76190 [*]	<.001

Table 3: Tukey post hoc test values



Bar graph 1: mean mortality for servo

2. Orchol leaf treatment

Normality test showed that data was normally distributed and was ready for the performance for further tests. Control values showed (SD=8.32, skewness 1.23±0.752, kurtosis=0.97±1.481). further normality test for 0.1%v/v concentration showed (SD=11.9, skewness 0.24±0.752, kurtosis=-1.6±1.481). Then normality for 0.5%v/v concentration showed (SD=6.39, skewness -.214±0.752, kurtosis=-.814±1.481). lastly for 1% values were (SD=10.06, skewness -1.525±0.752, kurtosis=3.481±1.481). One-way anova result showed

significant results. where sum of square between groups was 14325.377, df=3, mean square= 14325.377, f=47.345 with significance value <.001. Anova showed percentage mortality between groups sum of square as 2420.578, df= 24, mean square= 100.857. Total sum of square was 2420.578 with df=27. After performing post hoc test for orchol, it was found in table 5 that control, 0.1v/v, 0.5%v/v and 1% v/v are showing significant results. Mean difference between different treatments was highly significant. Average mean in table 6 for control means mortality in six days was 6.17% with (SD=8.33, range=23.33, std error= 2.94). 0.1%v/v orchol treatment showed average mean for mean mortality as 23.07% (SD=11.98, range=31.33, standard error of the mean=4.23. mineral oil with 0.5% v/v treatment gave aphid average mean mortality 50.74% with (SD= 6.44, range=18, Standard error of the mean=2.28. lastly 1%v/v orchol concentration showed aphid mortality in potato plants to 71.79% with (SD=10.07, range=33.33, and standard error of the mean= 3.55. bar graph showing increase in aphid mortality after orchol treatment with 0.1%v/v, 0.5%v/v and 1%v/v for six days. Maximum mortality was reported in 1%v/v spray treatment in aphids. Potato leaves didn't show any kind of variations in their colour after spray.

Percentage Mortality					
	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	14325.377	3	4775.126	47.345	<.001
Within Groups	2420.578	24	100.857		
Total	16745.955	27			


Table 4: showing one-way Anova result for servo concentrations

Type of treatment (I)	Type of treatment (J)	Mean difference (I-J)	significance
Control	0.1% orchol	-16.195*	<.001
	0.5% orchol	-43.742*	<.001
	1% orchol	-64.950*	<.001
0.1% Mortality	Control	16.195*	<.001
	0.5% orchol	-27.547*	<.001
	1% orchol	-48.755*	<.001
0.5% Mortality	Control	43.742*	<.001
	0.1% orchol	27.547*	<.001
	1% orchol	-21.207*	<.001
1% Mortality	Control	64.950*	<.001
	0.1% orchol	48.755*	<.001
	0.5% orchol	21.207*	<.001

Table 5: Tukey post hoc test values for orchol

	control mean mortality	0.1% mean mortality	0.5% mean mortality	1% mean mortality
Mean	6.9167	23.0716	50.7471	71.7917
Std. Deviation	8.33190	11.97084	6.44835	10.06162
Range	23.33	31.33	18.00	33.33
Std. Error of Mean	2.94577	4.23233	2.27984	3.55732

Table 6: Mean values for the average mean mortality in aphid population



Bar graph 2: graph showing mean for the average mean mortality in aphids after orchol treatments in potato plants.

3. Agrospele leaf treatment

Potato leaves were sprayed with 0.1% v/v, 0.5% v/v and 1% v/v mineral oil with trade name agrospele. Then aphids were allowed to feed for six days. Readings were observed on every day and aphid mortality in treatments were compared with control. Before proceeding to one-way anova normality test was done for the mean mortality readings. Normality test for average mean mortality in aphids after agrospele treatment showed skewness for control as 1.68 ± 0.75 and kurtosis to 2.81 ± 1.48 . Skewness and kurtosis for 0.1% v/v treatment is -1.35 ± 0.75 and 1.22 ± 1.48 . Treatment with 0.5% agrospele for aphid mortality gave skewness and kurtosis values as -1.88 ± 0.75 and 4.18 ± 1.48 . Anova results showed sum of square between groups to 21978.819, $df=3$, mean square=7326.273, $f=124.011$ with significance $<.001$. Sum of squares within groups is 1654.167 in table 7 with $df=28$ and mean square=59.077. Total sum of square is equal to 23632.986 with $df=31$ showing that our mean values are highly significant. Average mean in table 9 for control means mortality in six days was 5.00% with (SD=6.90, std error=2.43). 0.1% v/v agrospele treatment showed average mean for mean mortality as 20.4% (SD=8.62582, standard error of the mean=3.04969. mineral oil with 0.5% v/v treatment gave aphid average mean mortality 48.3% with (SD=8.35, Standard error of the mean=2.95. lastly 1% v/v agrospele concentration showed aphid mortality in potato plants to 73.3% with (SD=6.67, and standard error of the mean=2.35. Average mean in table 9 for control means mortality in six days was 5.00% with (SD=6.90, std error=2.43). 0.1% v/v agrospele treatment showed average mean for mean mortality as 20.4% (SD=8.62582, standard error of the mean=3.04969. mineral oil with 0.5% v/v treatment gave aphid average mean mortality 48.3% with (SD=8.35, Standard error of the mean=2.95. lastly 1% v/v agrospele concentration showed aphid mortality in potato plants to 73.3% with (SD=6.67, and standard error of the mean=2.35. Bar graph 3 showing increased trend in the mean of average mean mortality in aphids from lower to higher concentration of mineral oil spray used.

	Mean \pm SE	SD	Skewness	kurtosis
Control	5.00 \pm 2.43	6.86	1.68 \pm 0.75	2.81 \pm 1.48
0.1% agrospele	20.4 \pm 3.04	8.25	-1.35 \pm 0.75	1.22 \pm 1.48
0.5% agrospele	48.3 \pm 2.95	8.35	-1.88 \pm 0.75	4.18 \pm 1.48
1.0% agrospele	73.3 \pm 2.35	6.66	-1.14 \pm 0.75	1.35 \pm 1.48

Table 7: Normality readings for servo leaf treatment

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	21978.819	3	7326.273	124.011	<.001

UNDER PEER REVIEW