

Fecundity studies on near threatened Indian Butter catfish *Ompok bimaculatus* (Bloch, 1794) from Vamsadhara River, Andhra Pradesh

ABSTRACT

Ompok bimaculatus (Bloch 1794) is a near threatened catfish (IUCN, 2025), of freshwaters, belongs to the family Siluridae of the order Siluriformes. Due to its rich lipo-protein content and soft bony structure it becomes delicious and nutritiously highly valuable food fish species. At present *Ompok bimaculatus* facing decline in the natural populations, due to several anthropogenic factors such as indiscriminate fishing during the breeding season, unscientific use of pesticide in agricultural fields, gradual siltation in the freshwater habitat, depletion of water velocity as a result of decrease in water volume etc. Because of such drastic reduction in its population and a very restricted distribution in the freshwater ecosystem, Present work aims to an experimental analysis of reproductive biology of *Ompok bimaculatus*. Standardized knowledge on factors like fecundity, ova diameter, maturity, sex-ratio, and spawning are essential pre-requisites in fishery management and conservation. For understanding the dynamics of the gonads and to assess reproductive performance of species information of the phase of gonad development is virtually important.

Keywords: Food fish, Maturity, Gonado somatic Index, fecundity, spawning,

Introduction

The fresh water fish fauna of India is 3523 fish species belonging to 1097 genera and 272 families of 55 orders are recorded as per recent studies of ZSI, comprising into three main families known as, Cyprinidae, Siluridae and Channidae, inhabiting in the inland ecosystems (Kosygin *et al* 2024). However, majority of fishes are yet to be analysed well. Growth of the fish, length and weight parameters, have positive influence on reproduction. In general, the fecundity is more closely related to length than the somatic weight or ovary weight. Fecundity is elucidated in simple terms as number of potential eggs in the ovary. The discharge of eggs is known by a process called 'spawning'. Studies on the reproductive biology of fish is essential in evaluating the commercial potentialities of its stock, life history, cultural practice and

management of small fish's species (Eyo *et al.* 2014). Reproductive biology i.e., fecundity, spawning, sex ratio, etc. are among the import aspects of the biology of fishes which must be understood to explain the variations in the level of populations as well as to make efforts to increase the amount of yield. (K. Arambam *et al.* 2020; Gupta, *et al.* 2014).

Materials and methods

40 female individuals of *Ompok bimaculatus* species were collected randomly throughout the year from Mahendratana a major tributary of the Vamsadhara River, and local markets, and its originating from Mahendragiri Hills in the Gajapati district of Odisha and joining the Vamsadhara river upstream at Gotta barrage, in Andhra Pradesh. Collected fish weight and length were taken at the sampling site, afterward preserved in 4% formalin, further transferred and preserved in 10% formalin for laboratory studies.

Maturity stage

Maturity stages were dispensed based on the degree of development of testes in males and ovaries in females. Cycle of gonadal maturation has been studied by macroscopic and microscopic monthly examination of the different maturation stages of gonad; male and female gonads have then been grouped into different gonadal stages of development according to Nikolsky (1963).

Gonado-Somatic Index (GSI %)

The gonads expand to mature as the fish grow. Till the maturity reaches the ripened stage, the relationship of growth of gonads with that of fish is directly proportionate. However, with spawning, as the ova are delivered and the gonad is spent, the weight of it is likely to decrease. Hence, the GSI steadily increases till the development of gonads into ripening. Then there is a noteworthy decline with spawning. This index is calculated for both males and females separately and the monthly mean value was then plotted. Gonado-somatic index (GSI) was determined following the equation of Parameswarn *et al.* (1974):

$$\text{GSI} = (\text{Weight of gonad} / \text{Weight of fish}) \times 100.$$

Fecundity

Gravimetrically the fecundity of the fish was observed. The external connective tissues were detached carefully from the surface of the ovaries. With the help of blotting paper moisture of the ovaries was removed. Weight of the ovaries was recorded with fine electronic balance. Then 0.01 g of each ovary was taken out separately from anterior, middle and posterior regions of each ovarian lobe. The number of mature and immature eggs for each portion was sorted out and counted. The fecundity of the studied fish was observed adopting the methodology of LeCren (1951):

Fecundity = (number of eggs in the sample x weight of gonad) / weight of the sample.

Relationship of fecundity with various body dimensions such as body length, body weight, ovary length and ovary weight were transferred into log transformed and least squares regression equation (Bagenal 1978):

Ova diameter

At different stages of the maturity the diameter of the ovum was measured with ocular micrometer under stereoscopic Olympus microscope along the longest axis of the ovum (Clark 1934).

Results

Determination of Maturity stage

Four maturity stages of ovaries have been recognized on basis of microscopic and histological studies which are stated below:

Stage I (Immature): Ovaries translucent, colorless and thread like in appearance. Ova are not visible to naked eyes, but under microscope ova are irregular in shape, with a clear nucleus and transparent as yet yolk is not formed (Fig.2).

Stage II (Maturing): Ovaries yellowish white in colour and still thread like in appearance; ova are visible to naked eyes; under microscope ova are spherical in shape, partly opaque in appearance due to commencement of yolk development.

Stage III (Mature): Ovaries light yellowish in colour and enlarged in size; ova are clearly visible to naked eyes; under microscope spherical in shape and opaque in appearance except the transparent periphery (Fig. 1).

Stage IV (Ripe): Ovaries deep yellow in colour; with maximum size. Under microscope, ova are spherical in shape and opaque due to huge amount of yolk present. In this stage, ova are with their full size and start to become liberated through oviducts on putting light pressure on the abdomen.

Stage V (Spent): Ovaries pale whitish in colour; almost thread like in appearance resembling the Stage I ovary; distinguishable only with the presence of the pigmented covering of the ovary. Under microscope, ova are visible, very small in size, irregular in shape.

Immature gonads (Stage I) have been observed from March to May; highest percentage being observed in March while lowest percentage in May. Stage II (maturing gonad) first have been observed in May and available till September; highest percentage being observed in July and lowest percentage in September. Stage III (mature gonad) have been observed from March to October; highest percentage being observed in September and lowest percentage in March.

Ripe (Stage IV) have been observed from April to December with highest percentage being observed in November and lowest percentage in April. Spent (Stage V) have been observed in October to December; highest percentage being observed in December and lowest percentage in October.

Gonado-Somatic Index (GSI %)

The GSI value was low during September to April and increased monthly and attained its maximum level in June. In the female there were very low GSI (%) as observed during January to March. Thus, it represented the occurrence of pre-spawning period during January to March (Table:1, Fig.3).

The GSI value increased gradually to reach at the greater value in May and June (Spawning period), which was later on declined to minimum value.

Hence, it indicated that the pre-spawning period was developed during January to March. The GSI value enhanced gradually to a higher value in June (Spawning period) and then declined to minimum value in September (Table:1).

Fecundity

The fecundity of the studied fish individual varied from 3724 to 41552 eggs/fish. The mean value of eggs per gram of ovary varied in month from a minimum of 152 eggs/gm in September to a maximum of 2267 eggs/gm in July. The monthly variation of fecundity was calculated as mean sample. Fecundity was high during May- July, which declined sharply during August October and remained constant till January. Therefore, the length and weight of the fish species under study were directly correlated with fecundity. Additionally, there was a clear correlation between fecundity and ovarian weight. (Table:1, Fig.4).

Ova diameter

The diameter of immature eggs ranged from 0.211 mm- 1.345 mm. While before spawning the ova diameter range from 0.548 - 0.634 mm. In mature eggs the diameter ranged from 0.727 mm- 1.345 mm. The frequency of occurrence of eggs at different diameter showed that immature eggs were found out during February - April and mature eggs in the month of May- July (Table:1, Fig.5). Since August the development of mature eggs decreased quantitatively.

Discussion

Ompok bimaculatus the females were quantitatively dominant over males (Banik *et al.* 2012). Taiwo and Aransiola (2001) also noticed similar observations in some other catfish species. However, Ham (1981) pointed out that in cat fishes such dominance by female was due to certain environmental conditions while Fagade *et al.* (1984) pointed out the phenomenon because of population regulation. Chakrabarty *et al.* (2007) pointed out that the sexual maturity of *Ompok bimaculatus* discloses that during stage of maturity of the female the length reaches to a minimum of 17.0 cm. The present observation also depicts that the females are relatively longer (Banik and Malla 2011). For understanding the dynamics of the gonads and to assess reproductive performance of species information of the phase of gonad development is virtually

important. In this species the stage of maturity is noticed with the volume of gonad. Variations in the form of gonad were found to be noticed during filli-form appearance of the immature stage to the development of mature lobular stage during the maturation process. The maturation process stops with the formation of wrinkle-shaped gonad after completion of spawning. At that period the gonad looks like dull and lobule. Lamai (1993) also observed similar findings on the reproductive cycle and gonad development stages in different fish species.

De Vlaming (1972) deliberated the GSI (%) in teleosts and has used this parameter as indicator of the spawning period in teleost species. Further, considered the use of GSI (%) in reproductive biology which is noticed to be more appropriate when associated with other indicators of the reproduction under macroscopic observations (DeMartin and Lau 1999). Dmitincko (1970) pointed out that the GSI for female was always higher than that for male probably due to greater weight of the ripe female gonad. The fluctuations of GSI value confirmed that *Ompok bimaculatus* began to spawn in June and continued till August. In the present studies fecundity is relatively smaller in comparison to the other catfishes. Fecundity of *Silurus triostegus* in Ataturk Dam Lake varied from 6800 to 120,300 (Oymak *et al.* 2001).

Variations in fecundity of fish species may be due to selectively different factors such as feeding, food abundance, species differentiation, nutritional resources etc. (Bagenal 1978). Peak season of fecundity of *Ompok* species coincides with the onset of rain fall and flood water during May to August (Vanderwaals 1974). Harding (1966) stated that most of the tropical fishes are adopted to breed on the rising flood conditions. Thus, allowing the juveniles to take full advantage of the flooded banks for smooth feeding purpose and thus protected from predation. The choice of a particular season in fishes for breeding is influenced by various factors among these are changes in water quality characteristics, inter specific interactions and occurrence of suitable spawning sites (Zaret 1980 and Ward and Samarakoon 1981). The present findings also reveal a significant correlation between fecundity and ovary weight. The diameter of ova during peak season is ranged from 0.727 mm- 1.345 mm in the present observations. Douglas (1979) also noticed similar findings in *C. auratus*. The stage of the ovary of the studied fish species during October to January shows that the spawning period is lacking during a period from October to January.

Conclusion

Present study concludes with the finding of fecundity studies of butter catfish *Ompak bimaculatus*, however this study was incomplete, but it will be helpful for further studies on reproductive biology of *Ompak bimaculatus* in both female and male species.

Ethical statement

The fish species collected through cast netting and studied in the present study are not protected under The Wildlife Protection Act, 1972 (Last amended in 2013), Government of India, All the guidelines on animal use and care were followed accordingly.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

References

- Arthi T., Nagarajan S., Sivakumar A. A. and Vijayaraman K. 2013. Reproductive biology of two fresh water fishes, *Ompok bimaculatus* and *O. malabaricus* of the river Amaravathy, Tamil Nadu, India. Biolife.: 1(2): -45-53.
- Bagenal, T. 1978. Aspects of fish fecundity. In Ecology of Fresh Water Fish Production. 2nd edn, (Shelby, D. G. ed.), Black well: Oxford, London. pp. 101
- Banik, S. and Malla, S. 2011. Habitat Mapping for *Ompok pabda* (Bloch) in Gomoti River of Tripura. The 10th Agricultural Science Congress at National Bureau of Fish Genetic Resources (ICAR), Lucknow, U.P., India during 10th to 12th February, 2011, pp. 388
- Banik, S., Goswami, P. and Malla, S. 2012. Studies on breeding physiology of *Ompok pabda* (Bloch, 1794) in Tripura. U. P. J. Zool., 32(1): 67-72.

Chakrabarty, N.M., Mondal, S.C. and Chakrabarty, P.P. 2007. Artificial breeding seed production and rearing of butter fish *Ompok pabda* – A significant mile stone in technology advancement Fishing Chimes, 26 (10): 134 – 136

Clark, F.N. 1934. Maturity of the California Sardine (*Sardinella caerulea*), determined by ova diameter measurement. Fish. Bull. Sacramento, 42: 1-49.

De Martini, E.E. and Lau, B.B. 1999. Morphometric criteria for estimating sexual maturity in two snappers, *Etelis carbunculus* and *Pristopomoides sieboldii*. Journal of Fish Biology. 97: 449-458

De Vlaming, V.L. 1972. The effect of temperature and photoperiod on reproductive cycling in the estuarine gobiid fish (*Gillichthys mirabilis*). Fish, Bulletin. 73: 1137-1157.

Dmitincko, E.M. 1970. Reproduction of the sea catfish (*Arius thalassinus*: Rupel) in the Arabian. Res. J. Fish. and Hydrobiol. 3(1): 22-31.

Douglas, C. 1979. Sexual maturity and fecundity of African catfish, *Clarias gariepinus* with an observation on spawning behavior of the Nile catfish, (*Clarias lazera*), Zoological Journal of the Limnological society of London. 35: 1-45.

Eyo, V. O., Ekanem, A. P. and Jimmy U. U. 2014. A comparative study of the gonado-somatic index (GSI) and gonad gross morphology of African catfish (*Clarias gariepinus*) fed unical aqua feed and coppens commercial feed. Croatian J. Fisher. 72: 52-66.

Fagede, S.O., Adebisi, A.A. and Aatanda, A.N. 1984. The breeding cycle of tilapia, *Sarotherodon galilaeus* in the IITA Lake, Ibadan, Nigeria. Hydrobiologia. 100: 493-500.

Gupta, B. K., Sarkar, U. K. and Bhardwaj, S. K. 2014. Reproductive biology of Indian silurid catfish *Ompok pabda* in river Gumti. J. Environ. Biol. 35: 345 351.

Ham, R. 1981. The ecology of six native and two introduced fish species in Enoggera creek system, south east, Queens land. B.Sc (Hons) Thesis. Griffith Univ. Brisbane. pp. 157

Harding, D. 1966. Lake Kariba: the hydrology and development of fisheries. In Manmade lakes. 2nd edn. (R.H. Lowe Mc 25. Connel ed.). Academic press, London. pp. 123-134.)

K. Arambam, S.K. Singh, P. Biswas, A.B. Patel, A.K. Jena, P.K. Pandey, 2020. Influence of light intensity and photoperiod on embryonic development, survival and growth of threatened catfish, *Ompok bimaculatus* early larvae Journal of Fish Biology, 97 (2020), pp. 740-752

Kosygin, L., Mohapatra, A., Bineesh, K.K., Sharma, I., Jadhav, S.S., Khyndriam, D. (2024). Fauna of India Checklist: Pisces. Version 1.0. Zoological Survey India.

Lamai, S.L. 1993. Aspects of the applied biology of African catfish, *Clarias gariepinus* (Burchell, 1822). Aquacultural techniques and deltamethrin toxicity. Ph.D. Thesis, Dept. of Applied Zoology, University of Reading, U.K., pp. 204.

Le Cren, E.D. 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). J. Anim. Ecol. 20: 201-219.

Nikolsky, G.V. 1963. The Ecology of Fishes. Academic Press London, London, 352 pp

Oymak, S.A., Solak, K. And Unlu, E. 2001. Some biological characteristics of *Silurus triostegus* Heckel, 1843 from Ataturk Dam Lake (Turkey). Turkish J. Zool. 25 (2): 139-148.

Parameswarn, S., Sevaraj, C. and Radhakrishnan, S. 1974. Observation on the biology of *Labeo gonius* (Hamilton). Indian J. Fish. 21: 54-75.

Taiwo, I.O. and Aransiola, M.O. 2001. Length – weight relationship, condition factors and fecundity of two bagrid cat fishes, *Chrysich thysnigrodigitatus* and *Chrysichthys walkeri* in Asejire Lake. In Transactions of the Fisheries Society of Nigeria. Lagos. pp. 277. Vanderwaals, B.C.W. 1974. Observation on the breeding habits of African catfish *Clarias gariepinus* (Burchell). J. Fish Biol. 48: 6-10.

Ward J.A. and Samarakoon, J.I. 1981. Reproductive tactics of the Asian cichlids of the genus *Ectopoma* in Sri Lanka. Environ. Biol., 6: 95-103

Zaret, T.M. 1980. Life history and growth relationships of *Cichla ocellaris*, a predatory South American Cichlid. Biotropica, 12: 144-157

Table 1: Total length (cm), Total body weight of fish (g), weight of ovary (g), gonado-somatic index (GSI%), fecundity (Numbers), ova diameter (mm) of 40 individuals of *Ompok bimaculatus*

Months	Total Length (cm)	Total Weight (g)	Weight of Ovary (g)	Gonado-Somatic Index (GSI %)	Fecundity (Numbers)	Mean Ova Diameter (mm)
January	22.3	42.15	2.340	5.55	3724	0.211
	28.2	73.57	3.992	5.42	4213	0.327
	29.7	85.43	3.794	4.44	5698	0.243
February	26.4	88.20	3.435	3.89	4327	0.578
	27.2	74.50	4.232	5.68	3767	0.427
	28.5	90.20	4.878	5.40	3778	0.573
March	26.2	87.52	4.987	5.69	5520	0.544
	27.4	95.20	5.125	5.38	6549	0.634
	29.2	99.00	4.997	5.04	7289	0.765
April	25.4	88.20	4.237	4.80	7554	0.727
	26.3	85.75	4.759	5.54	8452	0.654
	24.7	68.07	5.880	8.63	8642	0.731
May	27.8	70.52	7.244	10.27	9239	0.765
	20.2	35.20	6.528	18.54	9768	0.711
	19.5	54.70	7.525	13.75	10880	0.755
June	24.7	70.50	8.721	12.37	12747	0.757
	20.5	38.00	8.280	21.78	12837	0.848
	34.7	105.50	10.280	9.74	15554	0.886
July	34.5	107.50	15.322	14.25	17121	1.102
	31.2	136.50	16.250	11.90	18009	1.107
	36.0	142.53	17.242	12.09	22552	0.928
August	32.4	145.24	16.827	11.58	24196	1.064
	31.0	115.00	13.180	11.46	27220	1.115
	28.7	107.20	12.412	11.57	23186	0.949
September	29.5	107.23	11.742	10.95	25597	0.756
	25.7	96.80	8.720	9.00	20157	0.812
	31.5	124.07	12.252	9.87	28525	1.345
October	32.7	127.15	12.774	10.04	29699	0.543
	24.8	77.50	6.235	8.04	15872	0.992
	20.2	44.50	6.335	14.23	13785	0.976
November	22.9	34.52	7.452	21.58	14293	1.042
	18.8	25.00	4.248	16.99	7934	0.875
	19.7	27.07	5.990	22.12	13781	0.720
December	18.4	23.50	4.087	17.39	9543	0.875
	20.5	28.50	5.209	18.27	11570	0.885
	21.7	32.72	4.528	13.82	9527	0.943
December	23.7	40.50	5.178	12.78	7190	0.925
	24.7	34.50	4.987	14.45	11192	0.942
	23.5	44.00	5.778	13.13	9934	0.875
	27.7	64.50	6.995	10.84	11123	1.175

(2 cm) of *Ompak*



Fig.1 Matured ovary
bimaculatus




Fig.2 Ovary weight of *Ompak bimaculatus*




Fig.3 Gonado-somatic Index of *Ompak bimaculatus*




Fig.4 Fecundity of *Ompak bimaculatus*




Fig.5 Ova diameter of *Ompak bimaculatus*