

# **Resistance Mechanisms in Aquatic Insects against Environmental Stressors: A Review**

## ABSTRACT

Aquatic insects are found nearly every kind of freshwater habitat and play an important role in aquatic ecosystem as predators, parasites, shredders, grazers, filter feeders, and piercers. Due to their critical role in energy flow and cycling of nutrients, they are essential components of freshwater ecosystems. However, their survival in a particular habitat is threatened by various environmental stressors including pollution, habitat degradation and climate change. These challenges affect their immune system thereby weakening their natural defenses. To fight against infections, aquatic insects primarily rely upon the innate immunity which includes both humoral and cellular responses. However, environmental factors such as fluctuating temperature, chemical pollutants and low oxygen level in aquatic body can damage these immune defenses, making the insects more susceptible to diseases. To defeat these challenges, some species have evolved adaptive strategies both physiological and behavioural. Such changes include the production of antimicrobial peptides, improved phenoloxidase activity and the generation of heat shock proteins. In the present review, we explore the existing research on how environmental stressors impact the immunity of aquatic insects. During our literature survey, we also examine the adaptive mechanisms the aquatic insects employ, the ecological implication of these changes and the importance of insect's immune adaptation in conservation efforts.

**Keywords:** *Aquatic insects, immunity, antimicrobial, conservation, ecological, adaptation*

## 1. INTRODUCTION

Insects belonging to the Phylum Arthropoda constitute the largest group not only of the animal kingdom but also of the whole living world. A large number of insects spend whole life or at least a part of their life in water called aquatic insects. Aquatic insects make up only 3-5 percent of all insect species, but are exceedingly taxonomically diverse (Daly *et al.* 1998). They occupy every kind of freshwater habitat including temporary streams and ponds, the shallowest and deepest areas of lakes, the most pristine and polluted rivers, roadside ditches, within and on macrophytes and susceptible to all ranges of water chemistry from acidified to alkaline bodies of water. They also represent all the functional feeding groups including predators, shredders, grazers (or scrapers), filter feeders, gatherers, piercers and parasites (Mackie, 2001). A study on aquatic insect fauna is one of the important aspects of freshwater biology. Aquatic insects form an important link in many food chains. They consume other invertebrates, small fish, aquatic plants, algae, detritus and decaying matter. Aquatic insects are also an important food source for birds, fishes, reptiles and amphibians. They have trophic link with terrestrial consumers like insectivorous birds (Danforth, 1926), which give them a higher consideration for the management of the wetland as wildlife refuge. Aquatic insects are also often used to determine water quality based on types and number of species. Many aquatic insects are biting pests to both humans and other animals. Some dipterans most notably mosquitoes are also vectors of diseases which include malaria, encephalitis and yellow fever. On the other hand adult odonates can greatly reduce adult mosquito population (Daly, *et al.* 1998) while the predaceous diving beetle (Coleoptera) and several other predatory aquatic insects (mostly Hemiptera) can help to reduce mosquito population by feeding on their larvae (Lundkvist *et al.*, 2003, Saha *et al.*, 2007). Midges feed on algal mats and sediments in sewage treatment facilities, which help them to keep them running properly (McCafferty, 1981). Some herbivorous aquatic insects have the potential to be used as biological control of invasive aquatic plants (Adler *et al.*, 2019).

The survival and evolutionary success of aquatic insects occupying various aquatic habitats are largely dependent on immune adaption and resistance mechanisms (Schmid-Hempel, 2005). They live in habitats that are abundant in bacteria, fungi, and viruses, as such they have evolved exceptional immune defences and infection-resistance mechanisms to survive in such pathogen-rich environments (Mahanta *et al.*, 2023). According to Flatt *et al.* (2008), aquatic insects mostly depend on their innate immune system, which consists of humoral and cellular reactions. Eleftherianos *et al.* (2021) found that hemocytes are essential for cellular immunity because they carry out processes like phagocytosis, encapsulation, and melanization. In contrast, the humoral response produces enzymes, other proteins, and antimicrobial peptides (AMPs) that destroy infections (Strand, 2008). The immunological responses of different insect species have changed as a result of adaptation to various aquatic habitats. While

some species strengthen their cuticle to prevent entrance of pathogens, others increase their ability to resist infections by producing more AMPs (Siva-Jothy *et al.*, 2005). Furthermore, immune memory-like responses have also been seen in insects despite their lack of vertebrate-like adaptive immunity, indicating that previous exposure to pathogens may improve subsequent immune responses (Cooper & Eleftherianos, 2017).

The molecular and cellular mechanisms of immunological adaptation in aquatic insects are still poorly understood, despite the comprehensive research on immune responses in model organisms (Little *et al.*, 2005; Rolff & Reynolds, 2009). The effects of pollution and temperature changes on aquatic organisms have been the focus of recent research, but little is known about how multiple stressors, including habitat loss, chemical exposure, and climate change, affect immune system function (Segner *et al.*, 2014). The aim of the present review is to scrutinize how immune system of aquatic insects living in different aquatic ecosystems is impacted by environmental stresses such as pollution, temperature swing, climate change, and habitat degradation. Additionally, it will assess the defence mechanisms that aquatic insects use against pathogen and parasite attacks.

## 2. ENVIRONMENTAL STRESSORS AFFECTING AQUATIC INSECT IMMUNITY

In a recent study carried out by Kataoka and Kashiwada, 2021, it has been clearly shown that environmental factors such as temperature change, oxygen depletion in aquatic environment and pollution play a major role on the defense mechanisms of aquatic insects, making them more vulnerable to illness. The same is also supported by Le Moullac (2000) with his study on environmental factors effecting the immune system of crustaceans, an important group aquatic arthropods. Increased water temperatures brought on by climate change can exacerbate thermal stress, which interferes with vital immune enzymatic functions (Siva-Jothy *et al.*, 2005). On the other hand, Vogelweith *et al.*, 2017 have found that long exposure to heavy metal lead and cadmium can damage the ability of an insect body to produce antimicrobial peptides which are critical elements to fight against infections. Environmental pollutants such as pesticides, heavy metals and industrial wastes also harm the defence mechanisms of aquatic insects reducing their ability to fight against diseases (Kim *et al.* 2021). This investigation highlights the immunotoxic effects on ecology, which can disrupt the stability of the ecosystem and population dynamics of aquatic insects. According to Rohr (2008), the key components of the innate immunity of aquatic insects such as haemocyte activity and phenoloxidase production are suppressed by the insecticides and pesticides used in agricultural activities. Eutrophication leading to reduced dissolved oxygen level cause hypoxic conditions in aquatic insects which potentially limit the energy resources necessary for the activation of the immune system (Ficke *et al.*, 2007; Brown *et al.* 2004). On the other hand, Gorr *et al.*, 2010 have already opined that hypoxia in aquatic environment disturbs the regulation of heat shock proteins (HSPs), preventing the process of cellular level apoptosis. Habitat destruction, one of the primary drivers of biodiversity change and sedimentation can alter the environment of aquatic habitats by affecting the availability of food resources and other important factors and indirectly affect immunity of aquatic insects by reducing energy intake and increasing stress (Dudgeon *et al.*, 2006; Townsend *et al.*, 2008).

In addition to these, pathogens such as bacteria, virus and fungi also affect the healthy life cycle of aquatic insects by altering their behaviour, reducing reproductive success and increasing mortality rates. For example, the fungus species *Lagenidium giganteum* causes high mortality in the mosquito population, especially in *Culex* (Merrim and Axtell, 1982; Jaronski and Axtell, 1983) and *Anopheles* species (Kerwin and Washino, 1987). Similarly the fungal pathogen *Metarhizium anisopliae* attacks the cuticle and internal tissues of mosquito larvae and reduces their survival rate (Scholte *et al.*, 2004). According to Williams *et al.* (2005), viral infections, such as *Iridovirus* cause mass mortality during larval stages in aquatic insects, which affects the food web dynamics and nutrient cycling. Understanding

these environmental impacts is cortically essential for conservation of aquatic ecosystems and reducing the effects of environmental change.




Figure 1. Diagrammatic representation showing Environmental Stressors and Immune Responses in aquatic insects

### 3. MECHANISMS OF IMMUNE ADAPTATION TO STRESSORS

The innate immune system in the form of phagocytosis, antimicrobial peptides and melanin-based encapsulation, is the most reliable form of cellular and humoral responses in aquatic insects (Siva-Jothy *et al.*, 2005). However, the immune systems of aquatic insects are continually challenged by the environmental stressors including pollution, temperature fluctuations and changes in water chemistry. The primary component of the aquatic insect immune system is the production of antimicrobial peptides (AMPs) which help in the protection against microbial infections by rupturing the cell membrane of the pathogens (Ezzati-Tabrizi *et al.*, 2013). The AMPs disorganize the microbial membranes by forming numerous pores causing cell lysis or interfering with the intracellular targets including DNA or protein synthesis (Bulet & Stocklin, 2005). Common AMPs identified in aquatic insect bodies include cecropins, effective against Gram-negative bacteria; defensins, which act against Gram-positive bacteria and attacins which target bacterial cell wall components (Vilcinskas, 2020). According to Cerenius & Soderhall (2004), the immune system of aquatic insects depends on phenoloxidase (PO) activity which is primarily involved in melanization, a process that encapsulates and neutralizes invasive foreign particles and microorganisms.

Pollutants like pesticides and heavy metals create major problems in aquatic insects by causing oxidative stress and interfering cellular signalling pathways and ultimately impair the immune system (Gill *et al.*, 2019). However, aquatic insects exhibit some remarkable adaptive changes through physiological and behavioural mechanisms. For example, some species demonstrate increased resistance to diseases by up regulating the PO activity (Laughton *et al.*, 2011). Furthermore, some species also produce heat shock proteins (HSPs), like HSP70 and HSP90, which serve as the molecular chaperones and shield the cellular proteins from thermal stress-induced damage (Pockley, 2003). Moreover, as indicated by Vilcinskas (2020), changes in the microbiome and epigenetic modifications such as DNA methylation and histone acetylation allows aquatic insects to adapt to changing environmental conditions, a key factor in immune plasticity.

### 4. ECOLOGICAL AND EVOLUTIONARY IMPLICATIONS

Environmental stressors, such as temperature fluctuations and pollution, significantly influence the immune adaptation of aquatic insects, with significant ecological and evolutionary consequences.

These stressors directly impact reproduction, survival rates, and population dynamics (Schmid-Hempel, 2011). According to Gonzalez-Santoyo and Cordoba-Aguilar (2012), immune adaptations influence species interactions, including prey-predator relationships and disease transmission. The recent review carried out by Schmid-Hempel (2005) on variation in immune defense as a question of evolutionary ecology have clearly opined that animal with stronger immune adaptations tend to live in pathogen-rich environments, outcompeting weaker ones, which leads to shifts in ecosystem dynamics and alterations in community composition. Cerenius and Soderhall (2004) also found that insects with higher levels of phenoloxidase activity and antimicrobial peptides (AMPs) are better equipped to resist pathogens compared to individuals lacking such adaptations leading to colonization in diverse habits. Such changes in immune system also influence reproductive success and population stability in aquatic insects as shown by Rolff and Siva-Jothy, 2003. Insects with weaker immunity expend more energy defending against the pathogens resulting in the fewer energy resources for reproduction and lower fecundity with slower population growth ( Rolff and Siva-Jothy, 2003). From the evolutionary point of view, long term exposure to environmental stressors can lead to natural selection, favoring species with stronger immune adaptations to survive and thereby passing their traits to next generations. Extreme environmental stressors however, exceed the ability of an organism's adaptive capacity which may lead to species extinction. This process may be a major driver of biodiversity change in aquatic ecosystems determining the species which can live in certain ecosystems and species which face population decline ( Rolff and Reynolds, 2009; Vilcinskas, 2020).

## **5. CONSERVATION PERSPECTIVES OF AQUATIC INSECTS**

Because of different environmental stressors, insects regularly face population crises, which can be solved by applying different conservation methods. A study by Burdon et al., (2019), discusses about the importance of conservation and the effects of pollutants on aquatic invertebrate communities. According to them, conservation should focus on building an environment-friendly home for insects, which includes reducing the use of chemical pollutants, maintaining their habitat integrity and supporting methods to reduce the consequences of climate change on insects. Studies about the immunological side of aquatic insects can help conservation planning by showing their level of adaptation to different stressors (Rolff and Reynolds, 2009). Habitat restoration and enhancing the water quality also help in reducing their complexity and improving the population size of the insects by strengthening their adaptive capacity (Palmer et al., 2010). According to Contador et al., (2012), the amount of research and publications about the conservation of aquatic insects is still very low. During their study, they found that from 1995 to 2008, only 5% of all the works published focused on the conservation of aquatic insects. More publications can attract more people's eyes towards aquatic insect conservation and their health-related issues.

## **6. CONCLUSION**

The present review highlights several key areas of research which need to be focused on. One major gap is the limited understanding of molecular and genetic mechanisms during immune adaptation in aquatic insect inhabiting different ecosystems. Recent advancement in the field of genetic and epigenetic studies of different organisms will be of key importance to identifying different pathways involved in immune responses to environmental stressors such as temperature change, pollution and hypoxia. Additionally, it has been found that a knowledge gap still exist regarding the combined examination of the effects of multiple stressors on aquatic insect populations. The role of micro-biota in regulating the immunity in insects is still unclear and further study is needed to clearly understand how environmental changes impact gut microbes of aquatic insects and effect immune functions.

## REFERENCES

- Adler, P. H., & Courtney, G. W. (2019). Ecological and societal services of aquatic Diptera. *Insects*, 10(3), 70. <https://doi.org/10.3390/insects10030070>
- Brown, G. C., Stock, N. E., & Chown, S. L. (2004). Oxygen supply limits the heat tolerance of insects. *Biological Reviews*, 79(2), 403–426. <https://doi.org/10.1017/S1464793103006355>
- Bullet, P., & Stocklin, R. (2005). Insect antimicrobial peptides: Structures, properties and gene regulation. *Protein & Peptide Letters*, 12(1), 3–11.
- Burdon, F. J., McIntosh, A. R., & Harding, J. S. (2019). Habitat loss drives threshold responses of benthic invertebrate communities to deposited sediment in agricultural streams. *Ecological Applications*, 29(2). <https://doi.org/10.1002/eap.1836>
- Cerenius, L., & Söderhäll, K. (2004). The prophenoloxidase-activating system in invertebrates. *Immunological Reviews*, 198(1), 116–126. <https://doi.org/10.1111/j.0105-2896.2004.00116.x>
- Contador, T. A., Kennedy, J. H., & Rozzi, R. (2012). The conservation status of southern South American aquatic insects in the literature. *Biodiversity and Conservation*, 21(8), 2095–2107. <https://doi.org/10.1007/s10531-012-0299-x>
- Cooper, D., & Eleftherianos, I. (2017). Memory and specificity in the insect immune system: Current perspective and future challenges. *Frontiers in Immunology*, 8, 539. <https://doi.org/10.3389/fimmu.2017.00539>
- Daly, H. V., Doyen, J. T., & Purcell, A. H. (1998). *Introduction to insect biology and diversity* (2nd ed.). Oxford University Press.
- Danforth, S. T. (1926). Birds of the Cartagena Lagoon, Porto Rico. *The Journal of the Department of Agriculture of Porto Rico*, 1, 137 pp.
- Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. I., Knowler, D. J., Lévêque, C., Naiman, R. J., Prieur-Richard, A. H., Soto, D., Stiassny, M. L. J., & Sullivan, C. A. (2006). *Freshwater*

- biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews*, 81(2), 163–182. <https://doi.org/10.1017/S1464793105006950>
- Eleftherianos, I., Heryanto, C., Basal, T., Zhang, W., Tettamanti, G., & Mohamed, A. (2021). Haemocyte-mediated immunity in insects: Cells, processes and associated components in the fight against pathogens and parasites. *Immunology*, 164(3), 401–432. <https://doi.org/10.1111/imm.13390>
- Ezzati-Tabarizi, R., Farookhi, N., Talaei-Hassanloui, R., Alavi, S. M., & Hosseininaveh, V. (2013). Insect inducible antimicrobial peptides and their applications. *Current Protein and Peptide Science*, 14, 620–631. <https://doi.org/10.2174/1389203711209070620>
- Ficke, A. D., Myrick, C. A., & Hansen, L. J. (2007). Potential impacts of global climate change on freshwater fisheries. *Reviews in Fish Biology and Fisheries*, 17(4), 581–613. <https://doi.org/10.1007/s11160-007-9059-5>
- Flatt, T., Heyland, A., Rus, F., Porpiglia, E., Sherlock, C., Yamamoto, R., Garbuzov, A., Palli, S. R., Tatar, M., & Silverman, N. (2008). Hormonal regulation of the humoral innate immune response in *Drosophila melanogaster*. *The Journal of Experimental Biology*, 211(Pt 16), 2712–2724. <https://doi.org/10.1242/jeb.01487>
- Gill, R., Garg, H., & Jain, S. (2019). Impact of environmental pollutions on insect immunity. *Environmental Toxicology and Pharmacology*, 68, 103226. <https://doi.org/10.1016/j.etap.2019.103226>
- Gonzalez-Santoyo, I., & Córdoba-Aguilar, A. (2012). Phenoloxidase: A key component of the insect immune system. *Entomologia Experimentalis et Applicata*, 142(1), 1–16. <https://doi.org/10.1111/j.1570-7458.2011.01187.x>
- Gorr, T., Cahn, J. D., Yamagata, H., & Bunn, H. F. (2010). Hypoxia-induced synthesis of hemoglobin in the crustacean *Daphnia magna* is hypoxia-inducible factor-dependent. *Journal of Biological Chemistry*, 285(40), 31767–31775. <https://doi.org/10.1074/jbc.M110.117002>
- Jaronski, S., & Axtell, R. C. (1983). Persistence of the mosquito fungal pathogen *Coelomomyces giganteum* (Oomycetes; Lagenidiales) after introduction into natural habitats. *Mosquito News*, 43, 332–337.
- Kataoka, C., & Kashiwada, S. (2021). Ecological risks due to immunotoxicological effects on aquatic organisms. *International Journal of Molecular Sciences*, 22(15), 8305. <https://doi.org/10.3390/ijms22158305>
- Kerwin, J. L., & Washino, R. K. (1987). Ground and aerial application of the asexual stage of *Lagenidium giganteum* for control of mosquitoes associated with rice culture in the Central Valley of California. *Journal of the American Mosquito Control Association*, 3, 59–64.
- Kim, T., Lee, S., & Choi, K. (2011). Ecological risks due to immunological effects on aquatic organisms. *International Journal of Environmental Research and Public Health*, 18(15). <https://doi.org/10.3390/ijerph18157837>
- Laughton, A. M., Fan, M. H., & Gerardo, N. M. (2011). The combined effects of bacterial symbionts and aging on immunity in a leaf-cutting ant. *Ecology and Evolution*, 1(4), 1–11. <https://doi.org/10.1002/ece3.3>
- Le Moullac, G. (2000). Environmental factors affecting immune responses in crustaceans. *Global Seafood Alliance*. <https://www.globalseafood.org>

- Little, T. J., Hultmark, D., & Read, A. F. (2005). Invertebrate immunity and the limits of mechanistic immunology. *Nature Immunology*, 6(7), 651–654. <https://doi.org/10.1038/ni1219>
- Lundkvist, E., Landin, J., Jackson, M., & Svensson, C. (2003). Diving beetles (Dytiscidae) as predators of mosquito larvae (Culicidae) in field experiments and laboratory tests of prey preference. *Bulletin of Entomological Research*, 93, 219–226.
- Mackie, G. L. (2001). *Applied aquatic ecosystem concepts*. Kendall/Hunt Publishing Company.
- Mahanta, D. K., Bhoi, T. K., Komal, J., Samal, I., Nikhil, R. M., & Paschapur, A. U. (2023). Insect-pathogen crosstalk and the cellular-molecular mechanisms of insect immunity: Uncovering the underlying signaling pathways and immune regulatory function of non-coding RNAs. *Frontiers in Immunology*, 14, 1169152. <https://doi.org/10.3389/fimmu.2023.1169152>
- McCafferty, W. P. (1981). *Aquatic entomology: The fisherman's and ecologist's illustrated guide to insects and their relatives*. Jones & Bartlett Learning.
- Merriam, T. L., & Axtell, R. C. (1982). Evaluation of the entomogenous fungi *Culicinomyces clavosporus* and *Coelomomyces giganteum* for control of the salt marsh mosquito, *Aedes taeniorhynchus*. *Mosquito News*, 42, 594–602.
- Palmer, M. A., Menninger, H. L., & Bernhardt, E. (2010). River restoration, habitat heterogeneity and biodiversity: A failure of theory or practice? *Freshwater Biology*, 55(s1), 205–222. <https://doi.org/10.1111/j.1365-2427.2009.02372.x>
- Pockley, A. G. (2003). Heat shock proteins as regulators of the immune response. *The Lancet*, 362(9382), 469–476. [https://doi.org/10.1016/S0140-6736\(03\)14075-5](https://doi.org/10.1016/S0140-6736(03)14075-5)
- Rohr, J. R. (2008). Agricultural insecticides in aquatic ecosystems. *Ecological Applications*, 18(4), 1011–1027. <https://doi.org/10.1890/07-1249.1>
- Rolff, J., & Reynolds, S. E. (2009). *Insect infection and immunity: Evolution, ecology, and mechanisms*. Oxford University Press.
- Rolff, J., & Siva-Jothy, M. T. (2003). Invertebrate ecological immunology. *Science*, 301(5632), 472–475. <https://doi.org/10.1126/science.1080623>
- Saha, N., Aditya, G., Bal, A., & Saha, G. K. (2007). A comparative study of predation of three aquatic heteropteran bugs on *Culex quinquefasciatus* larvae. *Limnology*, 8, 273–280.
- Schmid-Hempel, P. (2005). Evolutionary ecology of insect immune defenses. *Annual Review of Entomology*, 50(1), 529–551. <https://doi.org/10.1146/annurev.ento.50.071803.130420>
- Schmid-Hempel, P. (2011). *Evolutionary parasitology: The integrated study of infections, immunology, ecology, and genetics*. Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780199229482.001.0001>
- Scholte, E.-J., Knols, B. G. J., Samson, R. A., & Takken, W. (2004). Entomopathogenic fungi for mosquito control: A review. *Journal of Insect Science*, 4(19), 1–24. <https://doi.org/10.1673/031.004.1901>
- Segner, H., Schmitt-Jansen, M., & Sabater, S. (2014). Assessing the impact of multiple stressors on aquatic biota: The receptor's side matters. *Environmental Science & Technology*, 48(14), 7690–7696. <https://doi.org/10.1021/es405082t>
- Siva-Jothy, M. T., Moret, Y., & Rolff, J. (2005). Insect immunity: An evolutionary ecology perspective. In *Advances in Insect Physiology* (Vol. 32, pp. 1–48). Elsevier.

- Strand, M. R. (2008). The insect cellular immune response. *Insect Science*, 15(1), 1–14. <https://doi.org/10.1111/j.1744-7917.2008.00183.x>
- Townsend, C. R., Uhlmann, S. S., & Matthaei, C. D. (2008). Individual and combined responses of stream ecosystems to multiple stressors. *Journal of Applied Ecology*, 45(6), 1810–1819. <https://doi.org/10.1111/j.1365-2664.2008.01548.x>
- Vilcinskas, A. (2020). Epigenetic and microbiome-mediated immune adaptation in insects. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375(1809), 20190594. <https://doi.org/10.1098/rstb.2019.0594>
- Vogelweith, F., Moret, Y., Thiéry, D., & Moreau, J. (2017). Physiological mechanisms underlying the cost of immunity in insects. *Insect Science*, 24(4), 485–501.
- Williams, T., Barbosa-Solomieu, V., & Chinchar, V. G. (2005). A decade of advances in iridovirus research. *Advances in Virus Research*, 65, 173–248. [https://doi.org/10.1016/S0065-3527\(05\)65006-3-3](https://doi.org/10.1016/S0065-3527(05)65006-3-3)